



# Indiana University Health

IU Health Physicians Orthopedics & Sports Medicine

## **DENOVO (JUVENILE PARTICULATE CARTILAGE IMPLANTATION) OF FEMORAL CONDYLE**

### **PHYSICAL THERAPY PROTOCOL**

**Bryan M. Saltzman, M.D.**

Indiana University Health Physicians

Assistant Professor of Orthopaedic Surgery, Indiana University  
Sports Medicine, Cartilage Restoration, Shoulder/Elbow

IU Health Methodist Hospital – 1801 N Senate Ave, Indianapolis, IN 46202  
IU Health North – 201 Pennsylvania Pkwy #100, Carmel, IN 46280

317-944-9400

[www.bryansaltzmanmd.com](http://www.bryansaltzmanmd.com)

**Patient Name:** \_\_\_\_\_ **Date of Surgery:** \_\_\_\_\_

\_\_\_ **Evaluate and Treat**                      \_\_\_ **Provide patient with home program**

**Frequency:** \_\_\_\_\_ x/week x \_\_\_\_\_ weeks

	<b>WEIGHT BEARING</b>	<b>BRACE</b>	<b>ROM</b>	<b>EXERCISES</b>
<b>PHASE I</b> 0-6 weeks	Non-WB	<b>0-2 week:</b> Locked in full extension at all times	<b>0-6 weeks:</b> Use CPM for 6 hours/day, beginning at 0- 40°; advance 5- 10° daily as tolerated	<b>0-2 weeks:</b> Quad sets, SLR, calf pumps, passive leg hangs to 90° at home  <b>2-6 weeks:</b> PROM/AAROM to



		Off for CPM and exercise only  Discontinue after 2 wks		tolerance, patella and tibiofibular joint mobs, quad, hamstring, and glute sets, SLR, side-lying hip and core
<b>PHASE II</b> 6-8 weeks	Advance 25% weekly until full	None	Full	Advance Phase I exercises
<b>PHASE III</b> 8-12 weeks	Full	None	Full	Gait training, begin closed chain activities: wall sits, shuttle, mini-squats, toe raises  Begin unilateral stance activities, balance training
<b>PHASE IV</b> 12 weeks – 6 months	Full	None	Full	Advance Phase III exercises; maximize core/glutes, pelvic stability work, eccentric hamstrings  May advance to elliptical, bike, pool as tolerated
<b>PHASE V</b> 6-12 months	Full	None	Full	Advance functional activity  Return to sport-specific activity and impact when cleared by MD after 8 mos

**By signing this referral, I certify that I have examined this patient and physical therapy is medically necessary. This patient \_\_\_ would \_\_\_ would not benefit from social services.**

\_\_\_\_\_ **Date:** \_\_\_\_\_

**Bryan M. Saltzman, MD**