



Indiana University Health

IU Health Physicians Orthopedics & Sports Medicine

LATISSIMUS REPAIR

PHYSICAL THERAPY PROTOCOL

Bryan M. Saltzman, M.D.

Chief, Division of Sports Medicine & Shoulder/Elbow Surgery

Indiana University Health Physicians

Assistant Professor of Orthopaedic Surgery, Indiana University

Sports Medicine, Cartilage Restoration, Shoulder/Elbow Surgery

IU Health Methodist Medical Plaza North (MSK) – 201 Pennsylvania Pkwy #100,
Carmel, IN 46280

IU Health Methodist Hospital – 1801 N Senate Ave, Indianapolis, IN 46202
317-944-9400

www.bryansaltzmanmd.com

Patient Name: _____ **Date of Surgery:** _____

___ **Evaluate and Treat** ___ **Provide patient with home program**

Frequency: _____ x/week x _____ weeks

___ **Weeks 0-1:**

- Patient to do Home Exercises given post-op (pendulums, elbow ROM, wrist ROM, grip strengthening)
- Patient to remain in shoulder immobilizer for 6 weeks



Weeks 1-6:

- True PROM only! The tendon needs to heal back into the bone.
- ROM goals: 90° FF/30° ER at side; ABD max 40-60 without rotation
- No resisted motions of shoulder until 12 weeks post-op
- Grip strengthening
- No canes/pulleys until 6 weeks post-op, because these are active-assist exercises
- Heat before PT, ice after PT

Weeks 6-12:

- Begin AAROM → AROM as tolerated
- Goals: Same as above, but can increase as tolerated
- Light passive stretching at end ranges
- Begin scapular exercises, PRE's for large muscle groups (pecs, lats, etc)
- Isometrics with arm at side beginning at 8 weeks

Months 3-12:

- Advance to full ROM as tolerated with passive stretching at end ranges
- Advance strengthening as tolerated: isometrics → bands → light weights (1-5 lbs); 8-12 reps/2-3 sets per rotator cuff, deltoid, and scapular stabilizers
- Only do strengthening 3x/week to avoid rotator cuff tendonitis
- Begin eccentrically resisted motions, plyometrics (ex. Weighted ball toss), proprioception (es. body blade)
- Begin sports related rehab at 4 ½ months, including advanced conditioning
- Return to throwing at 4 months, begin with light toss
- Return to throwing from the pitchers mound at 6 months
- Return to full competition 9-12 months

Comments:

____ Functional Capacity Evaluation ____ Work Hardening/Work Conditioning ____ Teach HEP

Modalities

____ Electric Stimulation ____ Ultrasound ____ Iontophoresis ____ Phonophoresis ____ Heat
before/after ____ Ice before/after ____ Trigger points massage ____ TENS ____ Other
____ Therapist's discretion

By signing this referral, I certify that I have examined this patient and physical therapy is medically necessary. This patient ____ would ____ would not benefit from social services.



Date: _____

Bryan M. Saltzman, MD