



Indiana University Health

IU Health Physicians Orthopedics & Sports Medicine

OSTEOCHONDRAL ALLOGRAFT TRANSPLANTATION (OAT) WITH MENISCUS ALLOGRAFT TRANSPLANTATION (MAT) AND DISTAL FEMORAL OSTEOTOMY (DFO)

PHYSICAL THERAPY PROTOCOL

Bryan M. Saltzman, M.D.

Chief, Division of Sports Medicine & Shoulder/Elbow Surgery

Indiana University Health Physicians

Assistant Professor of Orthopaedic Surgery, Indiana University

Sports Medicine, Cartilage Restoration, Shoulder/Elbow Surgery

IU Health Methodist Medical Plaza North (MSK) – 201 Pennsylvania Pkwy #100,
Carmel, IN 46280

IU Health Methodist Hospital – 1801 N Senate Ave, Indianapolis, IN 46202

317-944-9400

www.bryansaltzmanmd.com

Patient Name: _____ **Date of Surgery:** _____

___ **Evaluate and Treat**

___ **Provide patient with home program**

Frequency: _____ x/week x _____ weeks

	WEIGHT BEARING	BRACE	ROM	EXERCISES
PHASE	Non-weight	Locked in full	Gentle passive 0-	Heel slides, quad sets, patellar



I 0-2 weeks	bearing	extension at all times* Off for hygiene and home exercise only	90° CPM 0-90°	mobs, SLR, calf pumps at home
PHASE II 2-8 weeks	2-6 weeks: Non-WB 6-8 weeks: Advance 25% weekly until full	2-6 weeks: Locked 0-90° Discontinue brace at 6 weeks	Advance as tolerated w/ caution during flexion >90° to protect post horn of meniscus	2-6 weeks: Add side-lying hip and core, advance quad set and stretching** 6-8 weeks: Addition of heel raises, total gym (closed chain), gait normalization, eccentric quads, eccentric hamstrings Advance core, glutes and pelvic stability
PHASE III 8-12 weeks	Full	None	Full	Progress closed chain activities Advance hamstring work, lunges/leg press 0-90° only, proprioception/balance exercises Begin stationary bike
PHASE IV 12-24 weeks	Full	None	Full	Progress Phase III exercises and functional activities: walking lunges, planks, bridges, swiss ball, half-bosu exercises Advance core/glutes and balance
PHASE V 6-9 months	Full	None	Full	Advance all activity w/o impact such as running, jumping, pivoting, sports until cleared by MD

*Brace may be removed for sleeping after first post-operative visit (day 7-14) **Avoid any tibial rotation for 8 weeks to protect meniscus

By signing this referral, I certify that I have examined this patient and physical therapy is medically necessary. This patient ___ would ___ would not benefit from social services.

Date: _____



Bryan M. Saltzman, MD